

# Bacterial and archaeal *amoA* gene distribution covaries with soil nitrification properties across a range of land uses

Lydia H. Zeglin,<sup>1\*</sup> Anne E. Taylor,<sup>1</sup> David D. Myrold<sup>1</sup> and Peter J. Bottomley<sup>1,2</sup>

Departments of <sup>1</sup>Crop and Soil Science and

<sup>2</sup>Microbiology, Oregon State University, Corvallis, OR, USA.

## Summary

Ammonia-oxidizing bacteria and ammonia-oxidizing archaea are commonly found together in soils, yet the factors influencing their relative distribution and activity remain unclear. We examined archaeal and bacterial *amoA* gene distribution, and used a novel bioassay to assess archaeal and bacterial contributions to nitrification potentials in soils spanning a range of land uses (forest, pasture, cultivated and long-term fallowed cropland) along a 10 km transect. The assay, which quantifies the extent to which acetylene-inactivated soil nitrification potential recovers (RNP) in the presence of bacterial protein synthesis inhibitors, indicated a significant archaeal contribution to the nitrification potentials of the pasture and long-term fallowed soils. Archaeal *amoA* gene abundance did not vary significantly among the soils, but bacterial *amoA* gene abundance did, resulting in archaeal : bacterial *amoA* abundance ratios ranging from  $1.1 \pm 0.8$  in cultivated soils to  $396 \pm 176$  in pasture soils. Both archaeal and bacterial *amoA* gene compositions were heterogeneous across the landscape, but differed in their patterns of variability. Archaeal *amoA* gene distributions were distinct among each of the three main land-use types: forest, pasture and cropland soils. In contrast, bacterial *amoA* gene composition was distinct in forest and in cultivated cropland, while pasture and long-term fallowed cropland soils were similar. In both pasture and long-term fallowed cropland soils, one phylotype of *Nitrosospira* cluster 3a was highly abundant. This distinct bacterial *amoA* gene fingerprint correlated with significant contributions of archaea to RNP of both

soils, despite differences in archaeal *amoA* gene composition between the pasture and fallowed soils. This observation suggests that the factors driving the development of ammonia-oxidizing bacteria community composition might influence the extent of archaeal contribution to soil nitrification.

## Introduction

Nitrification, the microbially mediated transformation of ammonium ( $\text{NH}_4^+$ ) to nitrate ( $\text{NO}_3^-$ ), is one of the most important components of the nitrogen (N) cycle. The rate-limiting step in this process is the oxidation of ammonia to nitrite ( $\text{NO}_2^-$ ). Ammonia oxidation by bacteria has been studied for many years, and takes place throughout terrestrial and aquatic environments. Recently, thaumarchaeal isolates from marine (*Candidatus Nitrosopumilus maritimus*) and soil (*Candidatus Nitrosopumilus viennensis*) have been shown to grow lithotrophically via ammonia oxidation in culture (Konneke *et al.*, 2005; Tournai *et al.*, 2011). Furthermore, several culture-independent soil studies have detected expression of the archaeal ammonia monooxygenase gene (*amoA*), suggesting that soil-borne archaea are capable of ammonia oxidation *in situ* (Treusch *et al.*, 2005; Leininger *et al.*, 2006; Chen *et al.*, 2008; Nicol *et al.*, 2008; Mertens *et al.*, 2009). Whereas *amoA* genes from ammonia-oxidizing thaumarchaea (AOA) tend to be numerically dominant over ammonia-oxidizing bacteria (AOB) in marine environments (Santoro *et al.*, 2010), soil environments are more variable with the ratio between archaeal and bacterial *amoA* genes ranging between 1 and 1000 (Francis *et al.*, 2005; Beman and Francis, 2006; Leininger *et al.*, 2006; Coolen *et al.*, 2007; He *et al.*, 2007; Adair and Schwartz, 2008; Boyle-Yarwood *et al.*, 2008). Several studies have shown that population size and community structure of archaeal, as well as bacterial, *amoA* genes are related to environmental conditions such as pH, salinity or N fertilization history (He *et al.*, 2007; Hansel *et al.*, 2008; Mosier and Francis, 2008; Nicol *et al.*, 2008; Shen *et al.*, 2008; Hallin *et al.*, 2009; Schauss *et al.*, 2009); however, to our knowledge there have been no direct comparisons of both AOA and AOB *amoA* gene community composition, population

Received 5 January, 2011; revised 22 June, 2011; accepted 16 August, 2011. \*For correspondence. E-mail lydia.zeglin@oregonstate.edu; Tel. (+1) (541) 737 3425; Fax (+1) (541) 737 3479.

sizes and nitrification activities in different soil types with varied land-use histories.

The factors affecting AOA and AOB distribution and their activities in the environment may be different (Erguder *et al.*, 2009). Generally, *Nitrosospira spp.* are the most common type of AOB found in soils. Vegetation type (Mintie *et al.*, 2003; Hawkes *et al.*, 2005) or N fertilization level (Avrahami *et al.*, 2003; Horz *et al.*, 2004) can affect the relative abundance of specific *Nitrosospira* taxa, and distinct *Nitrosospira* clades have been associated with soils expressing different nitrification properties (Webster *et al.*, 2005). Archaeal *amoA* genes exhibit phylogenetic structure definable by predominant habitat type ('soil' or 'marine') (Prosser and Nicol, 2008), and soil AOA abundance and activity has been shown to depend on pH and ammonium availability (Nicol *et al.*, 2008; Di *et al.*, 2010). In addition to pH, AOA community structure may also vary with fertilization practice (He *et al.*, 2007; Tourna *et al.*, 2008) or vegetation type (Boyle-Yarwood *et al.*, 2008; Chen *et al.*, 2008). Although it is now clear that some members of the Group 1.1a Thaumarchaea can oxidize ammonia, fix CO<sub>2</sub>, and grow in soil (Offre *et al.*, 2009; Zhang *et al.*, 2010; Verhamme *et al.*, 2011), soil-borne AOB usually proliferate more extensively than AOA in response to large additions of exogenous ammonia- or urea-N fertilizer (Di *et al.*, 2009; Jia and Conrad, 2009; Mertens *et al.*, 2009; Pratscher *et al.*, 2011; Verhamme *et al.*, 2011). Based on a novel assay that measures the extent of recovery of nitrification potential (RNP) over a 24–48 h incubation in the presence of bacterial protein synthesis inhibitors after irreversibly inactivating ammonia monooxygenase with acetylene, AOB and AOA were implicated as the dominant contributors to the nitrification potential of cropped and pasture soils respectively (Taylor *et al.*, 2010). However, it remains unknown whether popu-

lation sizes, community composition or edaphic properties control the relative contribution of AOA or AOB to nitrification activity. In order to understand the controls over contributions of AOA and AOB to *in situ* soil nitrification rates, it is necessary to identify factors that link diversity, abundance and activity of the AOA and AOB that coexist in field soils.

In this study, we investigated archaeal and bacterial *amoA* gene abundance (using Q-PCR), community composition (using T-RFLP and clone libraries), and nitrification activity in soils representing a range of land-use types (coniferous and actinorhizal forest, improved and unimproved pasture, actively cultivated and long-term fallowed cropland) from the Willamette Valley, Oregon, USA. Our goal was to identify patterns of coexistence of archaeal and bacterial *amoA* genes in these soils, and to determine if any relationship existed between these patterns and the relative contributions of archaea and bacteria to the RNP. A better understanding of the field distribution of *amoA* genes and nitrification activity in relation to varied soil management strategies will help to predict the relative importance of bacterial and archaeal nitrification across the landscape.

## Results and discussion

### *Soil properties, nitrification potential and amoA gene abundance*

Soils spanned a narrow pH range (6.0–6.7), a fourfold range in total C, a twofold range in total N and a sixfold range in net N mineralization rates (Table 1). Nitrification potential rates (NP) were high and variable, and were highest in the annually urea-N-fertilized cropped soils and lowest in the pasture soils, although this difference was

**Table 1.** Soil characteristics, N cycling parameters, and *amoA* gene abundance and richness.

	pH	Total C		Total N		Nmin µg N g <sup>-1</sup>	NP soil d <sup>-1</sup>	RNP-kan <sup>†</sup> (ratio)	Arch- <i>amoA</i> <sup>†</sup>	Bact- <i>amoA</i> <sup>†</sup>	Arch : Bact- <i>amoA</i> <sup>†</sup>	Arch- <i>amoA</i> T-RFs	Bact- <i>amoA</i> T-RFs
		(%)	C : N	Copies × 10 <sup>6</sup> g <sup>-1</sup> soil									
F-F	6.5 (0.1)	5.3 <sup>a</sup> (0.2)	0.27 <sup>abc</sup> (0.01)	23.0 <sup>a</sup> (0.9)	2.2 <sup>b</sup> (0.1)	5.8 (0.4)	na	9.32 (6.5)	0.51 <sup>ab</sup> (0.2)	13.7 <sup>b</sup> (7.5)	5 <sup>ab</sup> (0.9)	11 <sup>ab</sup> (0.6)	
F-A	6.7 (0.1)	5.2 <sup>a</sup> (0.9)	0.28 <sup>ab</sup> (0.05)	21.6 <sup>a</sup> (0.2)	4.4 <sup>a</sup> (0.7)	7.0 (5.0)	0.0 <sup>c</sup>	6.30 (1.4)	1.85 <sup>a</sup> (0.9)	10.6 <sup>b</sup> (8.6)	3 <sup>b</sup> (0.3)	12 <sup>a</sup> (0.1)	
P-L	6.0 (0.05)	3.6 <sup>abc</sup> (0.4)	0.26 <sup>abc</sup> (0.02)	16.2 <sup>b</sup> (0.6)	2.0 <sup>b</sup> (0.2)	1.8 (0.6)	na	11.1 (10.7)	0.08 <sup>b</sup> (0.07)	74.2 <sup>ab</sup> (39)	8 <sup>a</sup> (0.3)	6 <sup>c</sup> (1.7)	
P-H	6.0 (0.3)	4.5 <sup>ab</sup> (0.8)	0.33 <sup>a</sup> (0.05)	15.8 <sup>bc</sup> (0.2)	1.5 <sup>b</sup> (0.4)	2.9 (2.5)	0.9 <sup>a</sup> (0.2)	41.73 (21.2)	0.35 <sup>ab</sup> (0.3)	396 <sup>a</sup> (176)	7 <sup>a</sup> (0.3)	9 <sup>abc</sup> (1.5)	
C-F	6.4 (0.1)	2.1 <sup>bc</sup> (0.05)	0.17 <sup>bc</sup> (0.005)	13.9 <sup>bc</sup> (0.1)	1.1 <sup>b</sup> (0.3)	4.6 (0.6)	0.5 <sup>b</sup> (0.05)	3.88 (2.7)	1.02 <sup>ab</sup> (0.5)	3.3 <sup>b</sup> (1.3)	6 <sup>ab</sup> (0.1)	7 <sup>bc</sup> (0.5)	
C-C	6.3 (0.2)	1.4 <sup>c</sup> (0.04)	0.12 <sup>c</sup> (0.003)	13.4 <sup>c</sup> (0.4)	0.68 <sup>b</sup> (0.2)	23.9 (15.6)	0.1 <sup>c</sup> (0.03)	0.92 (0.7)	0.82 <sup>ab</sup> (0.2)	1.1 <sup>b</sup> (0.8)	5 <sup>ab</sup> (1.0)	11 <sup>ab</sup> (0.1)	

<sup>†</sup>. Modified from Taylor *et al.* (2010); Data are mean (SE), and superscripts denote differences based on one-way ANOVA ( $P < 0.05$ , Tukey *post hoc*).


F-F, Douglas-fir forest; F-A, alder forest; P-L, lightly grazed pasture; P-H, heavily grazed pasture; C-F, fallowed cultivated; C-C, cropped cultivated; Nmin, 30 day mineralizable N; Arch, archaeal; Bact, bacterial; na, not available.

not statistically significant. Recovery of NP (RNP) in the presence of bacterial protein synthesis inhibitors after acetylene inactivation varied from 0% to 104%, and was not correlated to the initial NP. Bacterial *amoA* gene abundance ranged over 20-fold, archaeal *amoA* gene abundance ranged almost 50-fold and the archaeal : bacterial *amoA* ratio spanned two orders of magnitude. Differences in putative ammonia-oxidizer population sizes between land-use types were driven by bacterial *amoA* gene abundance ( $P = 0.023$ ) and archaeal : bacterial *amoA* ratio ( $P = 0.016$ ). Pasture soils had the lowest bacterial *amoA* gene abundance and highest archaeal : bacterial *amoA* ratio, while forest, cultivated and long-term fallowed soils had the lowest archaeal : bacterial *amoA* ratio (Tukey *post hoc* comparisons,  $P < 0.05$ ).

Despite considerable differences among land-use types in nitrification rates (NP, RNP) and bacterial and archaeal *amoA* gene abundances, there were no statistically significant relationships between NP, RNP and either bacterial, archaeal or bacterial + archaeal *amoA* gene abundance or the archaeal : bacterial *amoA* ratio. Perhaps this result is to be expected given the limited knowledge of links between environmental *amoA* gene distribution and ammonia oxidizer activity. For example, it is well known that *in situ* gross rates of nitrification in whole soils range between 10% and 60% of the NP, implying that a variable fraction of soil ammonia oxidizers are  $\text{NH}_4^+$  limited, starved or inactive in whole soil situations (Norton, 2008). There are a few examples of pure cultures of AOB that, after  $\text{NH}_4^+$  starvation for between 1 and 5 weeks, can produce  $\text{NO}_2^-$  immediately upon addition of  $\text{NH}_4^+$  (Johnstone and Jones, 1988; Bollmann *et al.*, 2005; Berube *et al.*, 2007). However, within heterogeneous soil populations of AOB and AOA, the factors controlling recovery of ammonia oxidation after starvation remain unknown.

#### Archaeal and bacterial *amoA* gene T-RF clustering among soils


An examination of the relative abundances of soil bacterial and archaeal *amoA* gene terminal restriction fragment (T-RF) revealed diverse, heterogeneous archaeal and bacterial *amoA* gene distributions within and among land-use types (Figs 1 and 2). Although there was some overlap in individual T-RF occurrence, distinct archaeal *amoA* gene compositions were found in forest, pasture and cropland (including the long-term fallowed) soils (Figs 1A and 2A; MRPP *post hoc*:  $P < 0.002$ ,  $T < -4.5$ ,  $A > 0.21$ ). In contrast, bacterial *amoA* gene composition was distinct in forest soils (Figs 1B and 2B; MRPP *post hoc*:  $P < 0.001$ ,  $T < -6.0$ ,  $A > 0.30$ ) and in cropped soils (MRPP:  $P = 0.010$ ,  $T = -3.4$ ,  $A = 0.43$ ), but was similar in both pasture and long-term fallowed soils ( $P = 0.131$ , MRPP:  $T = -1.0$ ,  $A = 0.06$ ). These results imply that



**Fig. 1.** Nonmetric multidimensional scaling ordinations of *amoA* gene T-RF relative abundance for (A) archaeal and (B) bacterial *amoA* gene communities. Archaeal model parameters are stress = 3.182, instability = 0.011; bacterial model parameters are stress = 2.409, instability < 0.001. Site abbreviations are F-F, Douglas-fir forest; F-A, alder forest; P-L, lightly grazed pasture; P-H, heavily grazed pasture; C-F, fallowed cultivated; C-C, cropped cultivated. Significant correlations between community composition and environmental metrics (see Table 1 for abbreviation definitions) are shown by the vector overlay ( $P < 0.005$  with either axis).

whereas bacterial *amoA* gene composition has shifted from that of a cultivated soil to that of a pasture soil since the cessation of active cultivation (no N fertilization or tillage), the archaeal *amoA* gene composition has not changed significantly.

To identify soil properties and N cycling variables that were related to *amoA* gene composition, vectors representing the strongest correlations between the NMDS distance matrix and other study metrics were plotted (Fig. 1). C : N and N mineralization rate were correlated with both archaeal and bacterial *amoA* gene community structure, suggesting a commonality between those factors: in fact, the forest soils drove this relationship, being highest in C : N and N mineralization, and having



**Fig. 2.** Bar graphs of individual T-RF abundances at each sampling site (mean of field replicates,  $n = 3$  for all except C-F:  $n = 2$  and C-C:  $n = 4$ ) for (A) archaeal and (B) bacterial *amoA* gene communities. Site abbreviations are F-F, Douglas-fir forest; F-A, alder forest; P-L, lightly grazed pasture; P-H, heavily grazed pasture; C-F, fallowed cultivated; C-C, cropped cultivated. T-RF abundance is scaled to relative abundance, with 100% set as the sum of all T-RF peak areas for the bacterial *amoA* gene and 50% set as the sum of all T-RF peak areas for each restriction enzyme on the archaeal *amoA* gene.

both unique bacterial and archaeal *amoA* gene compositions. On the other hand, only the bacterial *amoA* gene composition was correlated to RNP + kanamycin (the fraction of RNP that is assumed to be attributable to AOA).

#### Archaeal *amoA* gene T-RF and sequence composition

Specific archaeal taxa were found to make up large percentages of the T-RF total fluorescence in forest, pasture and cropland soils (Fig. 2A). For example, on average, Rsa570 comprised 85% of the forest soil T-RF total fluorescence, Msp295 and Rsa631 constituted 25% and 33%, respectively, of the pasture soil T-RF fluorescence, and Rsa296 and Rsa200 contributed 47% and 38%, respectively, of the cropland soil T-RF fluorescence. Although vegetation has been suggested to influence soil archaeal community composition via rhizosphere effects (Chen *et al.*, 2008; Bomberg and Timonen, 2009), the archaeal *amoA* gene composition in forest soils was similar in both Douglas-fir and red alder stands, which

comprise different rhizosphere habitats. Additionally, archaeal *amoA* gene composition from this study showed no taxonomic overlap with the composition of archaeal *amoA* genes from Douglas-fir and red alder stands located on an acidic (pH 4) andisol about 100 km distant (Boyle-Yarwood *et al.*, 2008). All soils in this study had similar pH; however, they differed a great deal in age, texture, mineralogy and parent material (Parsons *et al.*, 1970; Taylor *et al.*, 2010). Although it is difficult to constrain and identify any one soil physical property as a driver of microbial community composition, soil texture determines pore space distribution and water dynamics, which can affect dispersal patterns and substrate availability for both microbial cells and microbial predators. Given the patterns we observed, it seems that vegetation type *per se* may be less important than edaphic factors as a driver of differences in archaeal *amoA* distribution among soils from differing land-use type.

It is notable that although several archaeal *amoA* gene T-RFs and sequences were detected in the study

soils, none corresponded with the T-RFLP profile or phylogenetic placement of 'Candidatus *N. maritimus*' (Msp631/Rsa119), 'Candidatus *Nitrososphaera gargensis*' (Msp631/Rsa196) or 'Candidatus *N. viennensis*' (Msp196/Rsa200) (Figs 2A and S1). However, an interesting pattern emerges when comparing the results of this study with others of archaeal *amoA* gene phylogenetic composition. Unlike any sequences collected in this study, the dominant archaeal *amoA* sequences identified by Boyle-Yarwood and colleagues (2008) in an acidic forested andisol fell into the same phylogenetic clades (iv and v) that were described in a Scottish agricultural loamy sand soil [Fig. S1; (Nicol *et al.*, 2008)]. In the latter study, archaeal *amoA* genes from clades iv and v were not only more abundant, but were also expressed to greater levels in acid soil (pH 4.3) than in the same soil that had been limed to maintain pH 6.9 (Nicol *et al.*, 2008). In addition, the dominant archaeal *amoA* gene phylotype that dominated the pH 6 forest soils of our study, T-RF Rsa570, fell into the same phylogenetic clade (iii) as sequences from the same Scottish agricultural soil (Fig. S1) that were more strongly expressed at pH 6.9 than 4.3 (Nicol *et al.*, 2008). The complimentary results of these three studies clearly suggest that pH is a soil factor affecting the abundance of certain phylotypes of archaeal *amoA* genes, regardless of soil type, local climate, vegetation or land use.

#### Bacterial *amoA* gene T-RF and sequence composition

The long-term fallowed and pasture soils had a bacterial *amoA* gene composition dominated by one phylotype (T-RF Alu200, 56% and 78%, respectively, of total fluorescence), unlike the forested and cropped soils, which contained a more even distribution of taxa (Fig. 2B). In addition, ordination analysis revealed that the long-term fallowed and pasture soils displayed a greater archaeal contribution to the RNP (RNP + kanamycin; Fig. 1B) than the cropped soils. Therefore, we should consider the possibility of a link between the factors that result in the dominance of T-RF Alu200 in the bacterial *amoA* gene community and the detection of a significant archaeal contribution to the RNP.

Although we cannot determine how many years without cultivation activity passed before the apparent shift of the bacterial *amoA* gene composition took place, the long-term fallowed soils had not been tilled, cropped, or N-fertilized for 18 years at the time of sampling. In calcareous grassland soils in the Netherlands, Kowalchuk and colleagues (2000) showed that the AOB community composition shifted noticeably following cessation of N fertilization over time periods ranging between 10 and 24 years. These shifts corresponded with the time needed to see substantial declines in net N

mineralization rates, suggesting that the ammonia oxidizers in non-fertilized soils were better adapted to living with lower  $\text{NH}_4^+$  availability. The Alu200 phylotype is related to sequences from *Nitrosopira* Cluster 3a (Table S1, Fig. S1). Previous work from the UK showed that unimproved, unfertilized pasture soils tend to contain *Nitrosopira* Cluster 3a, and these soils showed a trend for a significant delay in nitrification response to added N (Webster *et al.*, 2005). Culture-based studies confirmed that growth of representative isolates of *Nitrosopira* Cluster 3a is inhibited by high  $\text{NH}_4^+$  concentrations (Tourna *et al.*, 2010). Perhaps the dominance of the Alu200 phylotype in pasture and long-term fallowed soils, and the delay of *Nitrosopira* Group 3a isolates to respond to added  $\text{NH}_4^+$  is somehow linked to the high archaeal contribution to the RNP of these soils. Clearly, further studies are needed to identify the specific factors controlling the relative contributions of archaea and bacteria to nitrification in the long-term fallowed and pasture soils.

The contrasting patterns of soil archaeal and bacterial activity and *amoA* abundance and distribution suggest that environmental factors that vary at a landscape scale may affect AOA differently than AOB. The concept of niche differentiation among AOB phylotypes has been established for several years, following observations that soil AOB phylotypic composition differs in response to N availability (Bollmann *et al.*, 2002; Avrahami *et al.*, 2003; Tourna *et al.*, 2008), plant community composition (Hawkes *et al.*, 2005) temperature (Avrahami *et al.*, 2003), soil moisture (Avrahami and Bohannon, 2007), oxygen tension (Kowalchuk *et al.*, 1998), ammonia uptake kinetics and/or inhibition (Webster *et al.*, 2005; Tourna *et al.*, 2010). The range of factors that may influence AOB composition reflect the many possible niches that may exist for individual AOB phylotypes. In addition, recent studies suggest that soil AOA communities have different response patterns than co-located AOB to pH and temperature changes (Nicol *et al.*, 2008; Tourna *et al.*, 2008). In this study, there were no significant differences in archaeal *amoA* abundance or composition between 'high' and 'low' N situations within any land-use type, suggesting that N supply was less important to AOA distribution than AOB. AOA abundance and diversity have shown no or inconsistent responses to N fertilization in other studies (He *et al.*, 2007; Shen *et al.*, 2008; Di *et al.*, 2009; Hallin *et al.*, 2009; Jia and Conrad, 2009; Schauss *et al.*, 2009; Wang *et al.*, 2009), supporting the hypothesis that landscape-scale factors might mediate and/or dominate the influence of N amendment on soil AOA composition. In contrast, bacterial *amoA* gene composition was quite different in long-term fallowed soils than in adjacent cultivated and N-fertilized soils, thus N fertilization remains a potential driver of AOB distribution. Most impor-

tantly, our study suggests that differences in distribution patterns of archaeal and bacterial *amoA* genes may have ramifications for each group's respective role in soil nitrification across the landscape. Future research should further dissect the role of soil properties and patterns of land use on the distribution and activity of terrestrial nitrifying microbes.

## Experimental procedures

### Sites and sampling approach

We established a 10 km transect that spans the major landforms in the Willamette Valley, Oregon, USA, and represents the dominant land uses (i.e. vegetation and land management practices) and soil types. Three locations were selected near Corvallis, Oregon, USA (44.7°N 123.3°W), on lands owned and managed by Oregon State University: McDonald Forest (forest = F), Soap Creek Ranch (pasture = P) and Hyslop Farm (cropland = C). At each location, we selected two sites with histories of higher or lower N inputs. Soil chemical and N cycling properties for all soils are reported in Table 1.

The forest is dominated by Douglas-fir (*Pseudotsuga menziesii*) interspersed with stands of actinorhizal red alder (*Alnus rubra*) along water courses; sample points were located at ~350 m elevation, in soils of primarily a Jory-Gelderman complex (basalt-derived Haplohumults). The pasture is grazed land in a mid-elevation stream valley (elevation ~125 m), with soils of a Dixonville-Gellatly-Witham complex (basalt-derived Argixerolls and Haploxerolls). The cultivated location in the Willamette River floodplain (elevation ~85 m) has been the site of cropping trials since 1929, and includes actively cropped cereal (wheat, barley) fields and adjacent long-term fallowed fields that have been uncultivated and unfertilized since 1990 and are colonized by fescue grass (planted 1999), volunteer grasses and forbs. The soil is classified as a Woodburn silt loam (glaciolacustrine-derived Argixerolls). All sites have a Mediterranean climate (cool wet winters, warm dry summers), receive an annual mean precipitation of 90–130 cm and have an annual mean temperature of 10–12°C.

Soil N properties also differ among the three land-use types, ranging from high to low amounts of net mineralizable N along the continuum of forest, pasture and cultivated soils (Table 1). In order to sample soils representing a range of N inputs within each land-use type, we established paired sampling plots at each location. From the forest, we sampled soil in three Douglas-fir stands (F-F) and three adjacent stands of N<sub>2</sub>-fixing red alder (F-A). At the pasture, we sampled soil from three lightly grazed, upslope pastures (P-L) and three heavily grazed and manured bottomland pastures (P-H). At the cultivated location, we collected soil from four cropped and fertilized fields (single application of 90–180 kg urea-N ha<sup>-1</sup> year<sup>-1</sup>), two planted in wheat and two planted in barley (cropped sites were combined for statistical analysis, C-C), and two long-term fallow fields that had not been cultivated or fertilized for 18 years (C-F). Thus, for all parameters measured, field replicates were collected and analysed separately [*n* = 3 except C-C (*n* = 4) and C-F (*n* = 2)].

Soils analysed for DNA, total C and N and pH were collected in late July 2008. Soils analysed for NP, RNP and N mineralization (N min) assays were collected in late July 2009. We collected soils from the same plots at the same time of year in order to gather supplemental data on soil properties under similar field conditions. Although summer climate is generally stable at the study sites, temporal heterogeneity may affect variability in the data. At each of the 18 field plots, 4–6 soil cores (2.5 cm diameter by 10 cm depth) were gathered from points across each plot and homogenized for further analysis. Soils were carried back to the lab on ice and immediately sieved (2 mm) and roots removed. Soils were stored at either 4°C (soil properties, NP, RNP) or -20°C (DNA) until analysis.

### Soil chemistry and nitrogen cycling parameters

Soil pH was measured using standard methods (Klute, 1994; Sparks, 1994). Total organic C and N were measured on a Leco CNS-2000 Macro Analyzer (St. Joseph, MI, USA). N mineralization (N min) was calculated as the amount of inorganic N (NH<sub>4</sub><sup>+</sup> + NO<sub>2</sub><sup>-</sup> + NO<sub>3</sub><sup>-</sup> extractable in 0.5 M KCl) accumulated during a 30 day incubation at 30°C and 60% water holding capacity. NP was measured using a modified shaken soil slurry method (Hart *et al.*, 1994) as the amount of NO<sub>2</sub><sup>-</sup> + NO<sub>3</sub><sup>-</sup> formed over a 24 h incubation period (Taylor *et al.*, 2010). To measure RNP, assay slurries were held in an acetylene atmosphere for 6 h to completely inactivate ammonia monooxygenase, degassed and aerated, then incubated similarly to the NP assays for 48 h. Inferred archaeal (non-bacterial) contribution to RNP was assessed by adding kanamycin at the beginning of the recovery phase (Taylor *et al.*, 2010).

### Nucleic acid analysis

DNA was extracted from 0.25–0.30 g dry soil using a MoBio PowerSoil (Carlsbad, CA, USA) extraction kit, and quantified using a NanoDrop ND-1000 UV-Vis Spectrophotometer (Wilmington, DE, USA). PCR of the *amoA* gene was performed in 50 µl reaction volumes using approximately 50 ng of template DNA, 1 × PCR buffer (Promega Colorless GoTaq Reaction Buffer, Madison, WI, USA), 0.2 mM dNTPs, 1.2 Units Promega GoTaq (Madison, WI, USA), 0.04–0.12% BSA, and 0.2 µM each of either the bacterial *amoA*\_1R and *amoA*\_2F primers (Rotthauwe *et al.*, 1997; Mintie *et al.*, 2003; Boyle-Yarwood *et al.*, 2008), or the archaeal Arch-*amoA*-F and Arch-*amoA*-R primers (Francis *et al.*, 2005; de la Torre *et al.*, 2008). Thermalcycler protocols followed existing programs for archaeal (Francis *et al.*, 2005; Boyle-Yarwood *et al.*, 2008; de la Torre *et al.*, 2008) and bacterial *amoA* PCR (Mintie *et al.*, 2003).

For T-RFLP assays, 5' end 6-FAM-labelled forward primers were used to produce PCR products, which were purified using a QIAquick PCR Purification Kit (QIAGEN, Valencia, CA, USA), and restricted with TaqI and AluI for the bacterial *amoA* gene analysis (Horz *et al.*, 2000; Mintie *et al.*, 2003), and with RsaI and MspI for the archaeal *amoA* gene analysis (Boyle-Yarwood *et al.*, 2008). The digests were purified and fragment lengths and relative abundances were analysed

using an ABI 3100 capillary sequencer and Genotyper 3.7 (Foster City, CA, USA). Technical replicates were analysed on a subset (33%) of samples to ensure profile reproducibility. For each sample, any fragments comprising < 3% relative abundance of total fluorescence were removed from subsequent analysis. The relative fluorescence abundances of unique terminal restriction fragments (T-RFs) were exported for further analysis.

Cloning and sequencing was performed on both combined archaeal and bacterial PCR products from all samples; i.e. two 96-well plate libraries were constructed, one for each gene. A ligation of each batch of PCR products was produced using the TOPO TA kit (Invitrogen, Carlsbad, CA, USA), and submitted to the Genome Center at Washington University (St Louis, MO, USA). 96 clones per library were sequenced on an ABI 3730 capillary sequencer (Foster City, CA, USA) using both M13F and R primers. Sequences were trimmed and culled to remove all data with a *phred* value of < 20. Using CodonCode software (Dedham, MA, USA), remaining sequences were aligned and assembled to 2 × coverage and operational taxonomic units (OTUs) for the archaeal and bacterial *amoA* gene were defined at a sequence identity level of 99%. The highest quality representative sequence of each OTU was exported for phylogenetic analysis.

Q-PCR of the archaeal and bacterial *amoA* genes was performed using the Brilliant II SYBR Green Master Mix (Stratagene, La Jolla, CA, USA) and an ABI 7500 Real Time PCR System (Foster City, CA, USA). We used ~10 ng template DNA, a 25 µl reaction volume, and published primers and thermalcycler protocols for both bacterial [*amoA*\_1R and *amoA*\_2F primers (Rotthauwe *et al.*, 1997; Mosier and Francis, 2008)] and archaeal [Arch-*amoA*F and Arch-*amoA*R primers (Francis *et al.*, 2005; Mosier and Francis, 2008)]. Standard curves were constructed with  $2.0 \times 10^1$  to  $2.0 \times 10^{-4}$  ng *Nitrosomonas europaea* genomic DNA (bacterial *amoA*, efficiency = 91–94%,  $R^2 = 0.996$ ) or  $1.25 \times 10^1$  to  $1.25 \times 10^{-3}$  ng of a TOPO plasmid containing a *Nitrosopumilus maritimus amoBAC* gene insert (archaeal *amoA*, efficiency = 95–100%,  $R^2 = 0.994$ – $0.995$ ). Each reaction was run in triplicate, and replicates were analysed for a subset (33%) of samples to ensure quantitative accuracy. *AmoA* gene abundances were standardized by the mass of DNA recovered per g dry soil and  $\log_{10}$  transformed before statistical analysis.

#### Statistical analysis

To test for differences in quantitative variables (soil characteristics, N min, NP, RNP, T-RF richness, archaeal and bacterial *amoA* gene abundances) between land-use type, we used ANOVA with Tukey's *post hoc* analysis. Ordinations of community composition were constructed based on the relative abundance of T-RFs at all sites: nonmetric multidimensional scaling (NMS) scores were calculated from Sørensen's distances, a robust distance measure for datasets with many null values (Legendre and Gallagher, 2001). Highly significant correlations between NMS scores and edaphic properties or N cycling parameters ( $P < 0.005$  with either axis) were graphed using vectors overlaid on each NMS plot. Multiple response permutation procedure (MRPP) was used to test for treatment differences (land use, N input) in com-

munity composition. All ANOVA and correlation analyses were performed using SPSS 11 for Mac, and all multivariate analyses (NMS, MRPP) were performed using PC-ORD (McCune and Mefford, 1999).

The archaeal and bacterial *amoA* gene clone library sequence data were compared with the known sequences in the NCBI library and analysed phylogenetically to determine their taxonomic affiliations. The representative sequence from each OTU was compared with the NCBI nr/nt database using BLAST, and the GenBank number and percent sequence similarity of the closest match was recorded. A phylogenetic analysis was performed on all representative sequences from this study plus selected sequences from known isolates and environmental clones using ARB (Ludwig *et al.*, 2004) and PAUP 4.0 (Swofford, 2003). Phylogenetic trees were built using the neighbour-joining algorithm in ARB, and confirmed by PAUP with the general time-reversible model of nucleotide evolution as recommended by ModelTest (Posada and Crandall, 1998). Trees were bootstrapped in PAUP by resampling 1000 times using stepwise addition and tree bisection replacement. Finally, all sequences included in the phylogenetic analysis were restricted *in silico* with the same enzymes used for T-RFLP analysis of soil amplicons using CodonCode, and these results were used to designate phyla represented by the T-RFs. Sequences are stored in GenBank with accession numbers HM113498–HM113519.

#### Acknowledgements

This research was funded by USDA CSREES 2007–35107–1835. Thanks to Alex Krupkin, Rockie Yarwood, Stephanie Yarwood, Laurel Kluber and Shawn Starckenburg for lab assistance and/or advice. Soil physical and chemical properties were analysed at the Central Analytical Laboratory, Oregon State University. The Center for Genome Research and Biocomputing at Oregon State University provided T-RFLP analysis and Q-PCR instrumentation.

#### References

- Adair, K.L., and Schwartz, E. (2008) Evidence that ammonia-oxidizing archaea are more abundant than ammonia-oxidizing bacteria in semiarid soils of northern Arizona, USA. *Microb Ecol* **56**: 420–426.
- Avrahami, S., and Bohannon, B.J.A. (2007) Response of *Nitrosospira sp.* strain AF-Like ammonia oxidizers to changes in temperature, soil moisture content, and fertilizer concentration. *Appl Environ Microbiol* **73**: 1166–1173.
- Avrahami, S., Liesack, W., and Conrad, R. (2003) Effects of temperature and fertilizer on activity and community structure of soil ammonia oxidizers. *Environ Microbiol* **5**: 691–705.
- Beman, J.M., and Francis, C.A. (2006) Diversity of ammonia-oxidizing archaea and bacteria in the sediments of a hyper-nutriented subtropical estuary: Bahia del Tobari, Mexico. *Appl Environ Microbiol* **72**: 7767–7777.
- Berube, P.M., Samudrala, R., and Stahl, D.A. (2007) Transcription of all *amoC* copies is associated with recovery of *Nitrosomonas europaea* from ammonia starvation. *J Bacteriol* **189**: 3935–3944.

- Bollmann, A., Bar-Gilissen, M.J., and Laanbroek, H.J. (2002) Growth at low ammonium concentrations and starvation response as potential factors involved in niche differentiation among ammonia-oxidizing bacteria. *Appl Environ Microbiol* **68**: 4751–4757.
- Bollmann, A., Schmidt, I., Saunders, A.M., and Nicolaisen, M.H. (2005) Influence of starvation on potential ammonia-oxidizing activity and *amoA* mRNA levels of *Nitrosospira briensis*. *Appl Environ Microbiol* **71**: 1276–1282.
- Bomberg, M., and Timonen, S. (2009) Effect of tree species and mycorrhizal colonization on the archaeal population of boreal forest rhizospheres. *Appl Environ Microbiol* **75**: 308–315.
- Boyle-Yarwood, S.A., Bottomley, P.J., and Myrold, D.D. (2008) Community composition of ammonia-oxidizing bacteria and archaea in soils under stands of red alder and Douglas-fir in Oregon. *Environ Microbiol* **10**: 2956–2965.
- Chen, X.P., Zhu, Y.G., Xia, Y., Shen, J.P., and He, J.Z. (2008) Ammonia-oxidizing archaea: important players in paddy rhizosphere soil? *Environ Microbiol* **10**: 1978–1987.
- Coolen, M.J.L., Abbas, B., van Bleijswijk, J., Hopmans, E.C., Kuypers, M.M.M., Wakeham, S.G., and Damste, J.S.S. (2007) Putative ammonia-oxidizing Crenarchaeota in suboxic waters of the Black Sea: a basin-wide ecological study using 16S ribosomal and functional genes and membrane lipids. *Environ Microbiol* **9**: 1001–1016.
- Di, H.J., Cameron, K.C., Shen, J.P., Winefield, C.S., O'Callaghan, M., Bowatte, S., and He, J.Z. (2009) Nitrification driven by bacteria and not archaea in nitrogen-rich grassland soils. *Nat Geosci* **2**: 621–624.
- Di, H.J., Cameron, K.C., Shen, J.P., Winefield, C.S., O'Callaghan, M., Bowatte, S., and He, J.Z. (2010) Ammonia-oxidizing bacteria and archaea grow under contrasting soil nitrogen conditions. *FEMS Microbiol Ecol* **72**: 386–394.
- Erguder, T.H., Boon, N., Wittebolle, L., Marzorati, M., and Verstraete, W. (2009) Environmental factors shaping the ecological niches of ammonia-oxidizing archaea. *FEMS Microbiol Rev* **33**: 855–869.
- Francis, C.A., Roberts, K.J., Beman, J.M., Santoro, A.E., and Oakley, B.B. (2005) Ubiquity and diversity of ammonia-oxidizing archaea in water columns and sediments of the ocean. *Proc Natl Acad Sci USA* **102**: 14683–14688.
- Hallin, S., Jones, C.M., Schloter, M., and Philippot, L. (2009) Relationship between N-cycling communities and ecosystem functioning in a 50-year-old fertilization experiment. *ISME J* **3**: 597–605.
- Hansel, C.M., Fendorf, S., Jardine, P.M., and Francis, C.A. (2008) Changes in bacterial and archaeal community structure and functional diversity along a geochemically variable soil profile. *Appl Environ Microbiol* **74**: 1620–1633.
- Hart, S.C., Stark, J.M., Davidson, E.A., and Firestone, M.K. (1994) Nitrogen mineralization, immobilization, and nitrification. In *Methods of Soil Analysis, Part 2 Microbiological and Biochemical Properties*. Weaver, R.W., Angle, S., Bottomley, P., Bezdicek, D., Smith, S., Tabatabai, A., and Wollum, A. (eds). Madison, NJ, USA: Soil Science Society of America, pp. 985–1018.
- Hawkes, C.V., Wren, I.F., Herman, D.J., and Firestone, M.K. (2005) Plant invasion alters nitrogen cycling by modifying the soil nitrifying community. *Ecol Lett* **8**: 976–985.
- He, J.Z., Shen, J.P., Zhang, L.M., Zhu, Y.G., Zheng, Y.M., Xu, M.G., and Di, H.J. (2007) Quantitative analyses of the abundance and composition of ammonia-oxidizing bacteria and ammonia-oxidizing archaea of a Chinese upland red soil under long-term fertilization practices. *Environ Microbiol* **9**: 3152–3152.
- Horz, H.P., Rotthauwe, J.H., Lukow, T., and Liesack, W. (2000) Identification of major subgroups of ammonia-oxidizing bacteria in environmental samples by T-RFLP analysis of *amoA* PCR products. *J Microbiol Methods* **39**: 197–204.
- Horz, H.P., Barbrook, A., Field, C.B., and Bohannan, B.J.M. (2004) Ammonia-oxidizing bacteria respond to multifactorial global change. *Proc Natl Acad Sci USA* **101**: 15136–15141.
- Jia, Z.J., and Conrad, R. (2009) Bacteria rather than Archaea dominate microbial ammonia oxidation in an agricultural soil. *Environ Microbiol* **11**: 1658–1671.
- Johnstone, B.H., and Jones, R.D. (1988) Recovery of a marine chemolithotrophic ammonium-oxidizing bacterium from long-term energy-source deprivation. *Can J Microbiol* **34**: 1347–1350.
- Klute, A. (1994) *Methods of Soil Analysis. Part 1. Physical and Mineralogical Methods*. Madison, WI, USA: Soil Science Society of America.
- Konneke, M., Bernhard, A.E., de la Torre, J.R., Walker, C.B., Waterbury, J.B., and Stahl, D.A. (2005) Isolation of an autotrophic ammonia-oxidizing marine archaeon. *Nature* **437**: 543–546.
- Kowalchuk, G.A., Bodelier, P.L.E., Heilig, G.H.J., Stephen, J.R., and Laanbroek, H.J. (1998) Community analysis of ammonia-oxidising bacteria, in relation to oxygen availability in soils and root-oxygenated sediments, using PCR, DGGE and oligonucleotide probe hybridisation. *FEMS Microbiol Ecol* **27**: 339–350.
- Kowalchuk, G.A., Stienstra, A.W., Heilig, G.H.J., Stephen, J.R., and Woldendorp, J.W. (2000) Changes in the community structure of ammonia-oxidizing bacteria during secondary succession of calcareous grasslands. *Environ Microbiol* **2**: 99–110.
- Legendre, P., and Gallagher, E.D. (2001) Ecologically meaningful transformations for ordination of species data. *Oecologia* **129**: 271–280.
- Leininger, S., Urich, T., Schloter, M., Schwark, L., Qi, J., Nicol, G.W., et al. (2006) Archaea predominate among ammonia-oxidizing prokaryotes in soils. *Nature* **442**: 806–809.
- Ludwig, W., Strunk, O., Westram, R., Richter, L., Meier, H., Yadhukumar, et al. (2004) ARB: a software environment for sequence data. *Nucleic Acids Res* **32**: 1363–1371.
- McCune, B., and Mefford, M.J. (1999) PC-ORD. Multivariate Analysis of Ecological Data. Version 4. MjM Software, Gleneden Beach, Oregon, USA.
- Mertens, J., Broos, K., Wakelin, S.A., Kowalchuk, G.A., Springael, D., and Smolders, E. (2009) Bacteria, not Archaea, restore nitrification in a zinc-contaminated soil. *ISME J* **3**: 916–923.
- Mintie, A.T., Heichen, R.S., Cromack, K., Myrold, D.D., and Bottomley, P.J. (2003) Ammonia-oxidizing bacteria along meadow-to-forest transects in the Oregon cascade mountains. *Appl Environ Microbiol* **69**: 3129–3136.
- Mosier, A.C., and Francis, C.A. (2008) Relative abundance and diversity of ammonia-oxidizing Archaea and Bacteria



- in the San Francisco Bay estuary. *Environ Microbiol* **10**: 3002–3016.
- Nicol, G.W., Leininger, S., Schleper, C., and Prosser, J.I. (2008) The influence of soil pH on the diversity, abundance and transcriptional activity of ammonia oxidizing archaea and bacteria. *Environ Microbiol* **10**: 2966–2978.
- Norton, J.M. (2008) Nitrification in agricultural soils. In *Nitrogen in Agricultural Systems*. Schepers, J.S., and Raun, W.R. (eds). Madison, WI, USA: American Society of Agronomy, pp. 173–199.
- Offre, P., Prosser, J.I., and Nicol, G.W. (2009) Growth of ammonia-oxidizing archaea in soil microcosms is inhibited by acetylene. *FEMS Microbiol Ecol* **70**: 99–108.
- Parsons, R.B., Balster, C.A., and Ness, A.O. (1970) Soil development and geomorphic surfaces, Willamette-Valley, Oregon. *Soil Sci Soc Am Proc* **34**: 485–491.
- Posada, D., and Crandall, K.A. (1998) Modeltest: testing the model of DNA substitution. *Bioinformatics* **14**: 817–818.
- Pratscher, J., Dumont, M.G., and Conrad, R. (2011) Ammonia oxidation coupled to CO<sub>2</sub> fixation by archaea and bacteria in an agricultural soil. *Proc Natl Acad Sci USA* **108**: 4170–4175.
- Prosser, J.I., and Nicol, G.W. (2008) Relative contributions of archaea and bacteria to aerobic ammonia oxidation in the environment. *Environ Microbiol* **10**: 2931–2941.
- Purkhold, U., Pommerening-Roser, A., Juretschko, S., Schmid, M.C., Koops, H.P., and Wagner, M. (2000) Phylogeny of all recognized species of ammonia oxidizers based on comparative 16S rRNA and *amoA* sequence analysis: Implications for molecular diversity surveys. *Appl Environ Microbiol* **66**: 5368–5382.
- Rotthauwe, J.-H., Witzel, K.-P., and Liesack, W. (1997) The ammonia monooxygenase structural gene *amoA* as a functional marker: molecular fine-scale analysis of natural ammonia-oxidizing populations. *Appl Environ Microbiol* **63**: 4704–4712.
- Santoro, A.E., Casciotti, K.L., and Francis, C.A. (2010) Activity, abundance and diversity of nitrifying archaea and bacteria in the central California Current. *Environ Microbiol* **12**: 1989–2006.
- Schauss, K., Focks, A., Leininger, S., Kotzerke, A., Heuer, H., Thiele-Bruhn, S., et al. (2009) Dynamics and functional relevance of ammonia-oxidizing archaea in two agricultural soils. *Environ Microbiol* **11**: 446–456.
- Shen, J.P., Zhang, L.M., Zhu, Y.G., Zhang, J.B., and He, J.Z. (2008) Abundance and composition of ammonia-oxidizing bacteria and ammonia-oxidizing archaea communities of an alkaline sandy loam. *Environ Microbiol* **10**: 1601–1611.
- Sparks, D.L. (1994) *Methods of Soil Analysis. Part 3. Chemical Methods*. Madison, WI, USA: Soil Science Society of America.
- Swofford, D.L. (2003) *PAUP: Phylogenetic Analysis Using Parsimony. Version 4*. Sunderland, MA, USA: Sinauer Associates.
- Taylor, A.E., Zeglin, L.H., Dooley, S., Myrold, D.D., and Bottomley, P.J. (2010) Evidence for different contributions of archaea and bacteria to the ammonia-oxidizing potential of diverse Oregon soils. *Appl Environ Microbiol* **76**: 7691–7698.
- de la Torre, J.R., Walker, C.B., Ingalls, A.E., Konneke, M., and Stahl, D.A. (2008) Cultivation of a thermophilic ammonia oxidizing archaeon synthesizing crenarchaeol. *Environ Microbiol* **10**: 810–818.
- Tourna, M., Freitag, T.E., Nicol, G.W., and Prosser, J.I. (2008) Growth, activity and temperature responses of ammonia-oxidizing archaea and bacteria in soil microcosms. *Environ Microbiol* **10**: 1357–1364.
- Tourna, M., Freitag, T.E., and Prosser, J.I. (2010) Stable isotope probing analysis of interactions between ammonia oxidizers. *Appl Environ Microbiol* **76**: 2468–2477.
- Tourna, M., Stieglmeier, M., Spang, A., Konneke, M., Schintlmeister, A., Urich, T., et al. (2011) *Nitrososphaera viennensis*, an ammonia oxidizing archaeon from soil. *Proc Natl Acad Sci USA* **108**: 8420–8425.
- Treusch, A.H., Leininger, S., Kletzin, A., Schuster, S.C., Klenk, H.P., and Schleper, C. (2005) Novel genes for nitrite reductase and Amo-related proteins indicate a role of uncultivated mesophilic crenarchaeota in nitrogen cycling. *Environ Microbiol* **7**: 1985–1995.
- Verhamme, D.T., Prosser, J.I., and Nicol, G.W. (2011) Ammonia concentration determines differential growth of ammonia-oxidising archaea and bacteria in soil microcosms. *ISME J* **5**: 1067–1071.
- Wang, Y.A., Ke, X.B., Wu, L.Q., and Lu, Y.H. (2009) Community composition of ammonia-oxidizing bacteria and archaea in rice field soil as affected by nitrogen fertilization. *Syst Appl Microbiol* **32**: 27–36.
- Webster, G., Embley, T.M., Freitag, T.E., Smith, Z., and Prosser, J.I. (2005) Links between ammonia oxidizer species composition, functional diversity and nitrification kinetics in grassland soils. *Environ Microbiol* **7**: 676–684.
- Zhang, L.-M., Offre, P.R., He, J.-Z., Verhamme, D.T., Nicol, G.W., and Prosser, J.I. (2010) Autotrophic ammonia oxidation by soil thaumarchaea. *Proc Natl Acad Sci USA* **107**: 17240–17245.

## Supporting information

Additional Supporting Information may be found in the online version of this article:

**Fig. S1.** Neighbour-joining phylogenetic tree of Thaumarchaeal *amoA* gene sequences from this study (99% OTU representatives marked in bold, number represents that OTU's abundance in the clone library), cultured isolates and other environmental sequences. *In silico* determinations of T-RFs for all sequences are noted to the right of the phylogram. T-RFs that represent monophyletic taxonomic groups are highlighted in bold/italic type, and the corresponding clade is marked with a unique symbol.

**Fig. S2.** Neighbour-joining phylogenetic tree of bacterial *amoA* gene sequences from this study (99% OTU representatives marked in bold, number represents that OTU's abundance in the clone library), characterized *Nitrosospira* sp. isolates and other environmental sequences. *In silico* determinations of T-RFs for all sequences are noted to the right of the phylogram. T-RFs that represent monophyletic taxonomic groups are highlighted in bold/italic type, and the corresponding clade is marked with a unique symbol.

**Table S1.** Archaeal and bacterial *amoA* gene clone library sequence information. AOB taxonomic identity based on a sequence grouping in a clade with known isolates (Fig. S2); cluster designation based on Purkhold and colleagues (2000).

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.